

# **Harlequin Duck Research and Monitoring in Montana: 1998**

A Report to:

ASARCO, Incorporated  
Box 868  
Troy, MT 59935

Submitted by

PAUL HENDRICKS

Montana Natural Heritage Program  
1515 East Sixth Avenue  
Helena, Montana 59620-1800

March 1999

Call #: S

639.97841

N11HDM

1999

1

Barcode:

Montana State Library



3 0864 1004 7140 1

00 000001

000000

© 1999 Montana Natural Heritage Program

This document should be cited as follows:

Hendricks, P. 1999. Harlequin Duck research and monitoring in Montana. 1998. Montana Natural Heritage Program. Helena, MT. 30 pp.

## **ABSTRACT**

In 1998, approximately 235 km of Harlequin Duck stream surveys were conducted on six streams, with the majority of effort focused on four Lower Clark Fork streams (Rock Creek, Swamp Creek, Marten Creek, Vermilion River). Additional streams surveyed were Trail Creek (North Fork Flathead River drainage) with one pair survey and one brood survey, and Upper McDonald Creek (Glacier National Park) with one brood survey.

On the Lower Clark Fork streams, stream flow was significantly below normal during early-May pair surveys. A minimum of 24 adult Harlequins (13 males, 11 females) was seen on 3 streams. These included Marten Creek (7 pairs, 2 males), Rock Creek (1 pair, 1 male, 1 female), Vermilion River (2 pairs) and Swamp Creek (no birds). The number of birds observed was about average for all streams, although Swamp Creek again had no birds.

Brood surveys were conducted during July and early August 1998. Seven broods were observed on the Lower Clark Fork streams (2 each on Marten Creek and Vermilion River, 3 on Rock Creek), successfully producing a maximum 11 young on Marten Creek, 6 young on Vermilion River, and 13 young on Rock Creek. Age of broods indicated that nesting among these streams was initiated over a period of at least three or four weeks.

New birds were banded on Marten Creek (2 males, 2 females, 5 juveniles), Vermilion River (1 female, 3 juveniles) and Rock Creek (2 juveniles). Additionally, several previously marked birds were re-observed, including 4 third-year females banded as juveniles in 1995 (3 of which successfully raised broods on their natal streams). One third-year female, banded as a juvenile on Swamp Creek in August 1995, was observed paired in May 1998 on Rock Creek, an inter-stream movement of ca. 21 km, and the first documented between these two streams.

Stream survey results from other areas were as follows. On Trail Creek (N. Fork Flathead drainage) 3 pairs and 1 male were counted on 14 May, 1 female (already banded) with a brood of 1 juvenile was captured and banded on 7 August. On McDonald Creek in Glacier National Park, park personnel reported a maximum of 8 pairs in spring. During brood surveys, 2 females (1 already banded) with 2 broods (8 juveniles) were captured and banded on 6 August. Three additional broods (3 females, 13 juveniles) were counted on 17 and 19 August by park personnel, and 2 females (1 already banded) and 12 juveniles were banded.

## **ACKNOWLEDGMENTS**

Much field assistance was provided by Bryce Maxell, who conducted most of the late-spring and early-summer surveys. Additional field help, identification of possible trapping sites, and other logistical support were provided by J. Ashley, especially in Glacier National Park. C. Jones, and M. Miller assisted with data entry and map preparation. J. Elliott served as a liaison to ASARCO, Inc., and reviewed an earlier draft of this report. Information from surveys conducted by other agencies was provided by John Ashley (Glacier National Park). Thanks to Ken Wright, Cyndi Smith, and Janet Hardin for sharing reports and sightings of marked ducks from the Pacific Coast. Primary financial support for the project came from ASARCO, Inc. (through the generosity of David Young), with additional support from Glacier National Park (thanks to Steve Gniadek) and the Montana Natural Heritage Program (Montana State Library, Natural Resource Information System and The Nature Conservancy cooperating).

## TABLE OF CONTENTS

ABSTRACT	iii
ACKNOWLEDGMENTS	iv
INTRODUCTION	1
METHODS AND MATERIALS	2
SURVEYS AND BANDING	3
MOVEMENT	6
DEMOGRAPHY AND POPULATIONS	9
CONSERVATION AND MANAGEMENT	12
PRIORITIES FOR FUTURE RESEARCH	13
REFERENCES	16
APPENDICES	28



## INTRODUCTION

The Harlequin Duck (*Histrionicus histrionicus*) is a small sea duck, which travels inland to breed on fresh water streams. Approximately 150-200 pairs of Harlequins currently breed in Montana (Reichel and Genter 1995), with most located in the following areas: 1) tributaries of the lower Clark Fork River, 2) tributaries of the North, Middle, and South Forks of the Flathead River, 3) streams coming off the east front of the Rocky Mountains; and 4) the Boulder River (Miller 1988, 1989, Kerr 1989, Carlson 1990, Fairman and Miller 1990, Diamond and Finnegan 1992, 1993, Reichel and Genter 1993, 1994, 1995, 1996).

During the breeding season, Harlequins are found along fast mountain streams (Bengtson 1966). In many areas, Harlequins use streams with dense timber or shrubs on the banks (Cassirer and Groves 1990), but they are also found in relatively open streams along the east slopes of the Rocky Mountains, Montana (Markum and Genter 1990, Diamond and Finnegan 1992), and in the Arctic tundra (Bengtson 1972). In Idaho, 90% of observations occurred near old growth or mature timber stands (Cassirer and Groves 1990). Mid-stream rocks, logs, islands, or stream-side gravel bars serve as safe loafing sites and appear to be important habitat components.

Most ducks arrive on their inland breeding areas in mid-April to early-May; unmated males typically arrive before pairs (Kuchel 1977). The males return to the coast shortly after the females begin incubation; most are gone by early July (Kuchel 1977). The females and young remain on the streams until August or early September. This chronology is influenced by elevation and by the timing of spring runoff; it may vary up to several weeks between years.

The Harlequin Duck is listed as "Sensitive" by the U.S. Forest Service, Region 1 (Reel *et al.* 1989) and as a Species of Special Concern by the Montana and Idaho Natural Heritage Programs (Idaho Conservation Data Center 1994, Montana Natural Heritage Program 1997).

The Montana Natural Heritage Program began banding Harlequin Ducks to a limited extent in 1991. Through 1997, a total of 356 Harlequin Ducks were marked on 9 streams, representing the largest marked "population" from the breeding streams. Birds marked in Montana have subsequently been captured and observed on the coasts of Oregon, Washington and British Columbia, with most reports coming from Vancouver Island (Reichel and Genter 1996, Reichel *et al.* 1997).

Objectives for the 1998 season included 1) surveying the Lower Clark Fork streams (Rock Creek, Marten Creek, Swamp Creek, Vermilion River) of Sanders County for presence and status of Harlequin Ducks, 2) gathering duck productivity data on the Lower Clark Fork streams, 3) marking as many individuals as possible on these streams for long-term monitoring, and 4) surveying McDonald Creek in Glacier National Park, the only other population in Montana with survey data comparable to the Lower Clark Fork population.

This report summarizes results from the 1998 field season and should not be considered a comprehensive analysis of all data collected from previous years (although some of the older data are presented). For a recent and relatively comprehensive summary of Harlequin Duck research in Montana through 1996 see Reichel *et al.* (1997). The 1997 field season is reported in Hendricks and Reichel (1998).

## **METHODS AND MATERIALS**

Harlequin Ducks were surveyed on parts of the Kootenai National Forest and in Glacier National Park during May-August 1997. Area of focus included the four principal breeding streams of the Lower Clark Fork population in Sanders County, Montana. These Clark Fork River tributaries are Rock Creek, Marten Creek (including the S. Fork Marten Creek), Swamp Creek, and Vermilion River. Single pair and brood surveys were also conducted on Trail Creek (North Fork Flathead River) in Flathead County, as this stream might function as a movement corridor into Glacier National Park. Surveys were conducted on McDonald Creek in Glacier National Park by Park personnel (pair surveys) and MTNHP personnel in conjunction with Park personnel (brood surveys).

Surveys were conducted by walking the stream channel (when possible) or stream bank. In most cases, the surveyor walked upstream, giving more time to observe the bird before it moved out of sight; in cases where birds were not to be marked, the surveyor made a loop around the birds to minimize disturbance. For streams in the Flathead and Clark Fork drainages, we attempted to capture and mark all birds seen when a licensed, qualified bird-bander was present on the survey.

Captured birds were sexed, aged, weighed, measured (wing cord and tail), marked, and released. Juveniles were aged based on feather development: Class IA-C: downy (1-14 days old), no feathers visible; Class IIA-C: partly feathered (15-35 days old); Class III: fully feathered but flightless (36-51 days old). Birds were banded with a USFWS aluminum band and with a blue, plastic leg band with 2 white alpha alpha (juveniles) or alpha numeric (adults) characters. These birds are individually recognizable by the imprinted characters, although the bands are less readily observed than nasal discs, which were also used in years prior to 1997 but discontinued because of concerns about trauma and pairing success. Dates, locations, distance surveyed, and general characteristics of the stream reaches surveyed were recorded; location, number, age, and sex of all Harlequin Ducks seen were recorded, as were habitat characteristics of the sites where ducks were first observed. All surveys and duck observations were entered into a database and associated ARC-INFO coverages.

Not all streams used by Harlequin Ducks during the breeding season are used for nesting or brood rearing. Some streams where adult Harlequins are observed may be used only during migration to and from breeding areas. In order to classify Harlequin Duck observations in a consistent manner MTNHP has adopted the following definitions proposed by Cassirer et al. (1996) (the first two of which would be considered "Element Occurrences" [EOs] by Natural Heritage Programs/Conservation Data Centers throughout North America). See previous reports (Reichel and Genter 1996, Reichel et al. 1997) for a full discussion of the distribution of Harlequin Ducks throughout Montana, and the status and classification for each occupied stream or drainage.

### **Harlequin Duck breeding occurrence:**

is defined by a drainage, drainages, or portion of a drainage where breeding is known (i.e., a brood or nest has been observed within the last 15 years).

EOs are separated by either:

- A substantial barrier (>2 km over a major divide); or,
- A 10-km separation for completely unsuitable habitat (across land);
- A 20-km separation (measured along watercourses) for both rarely used habitat (lakes, <1% gradient rivers) and for apparently suitable habitat



that is not known to be occupied

**Probable harlequin duck breeding occurrence:**

Same definition as above, except breeding is not known, but rather is highly suspected (i.e., there have been at least 3 independent pair or female observations within the last 15 years)

**Breeding status unknown:**

Drainages or portions of drainages with at least 1 harlequin duck observation but fewer than 3 independent pair or female observations during the breeding season within the last 15 years

**Breeding unlikely:**

Observations of males during migration periods. The male migration periods are before 15 April and after 5 June in the Northern Columbia Basin and Rocky Mountain Front areas and before 1 May and after 20 June in the Intermountain region

Observations of pairs outside the pre-nesting season. The pre-nesting season is from 15 April - 5 June in the Northern Columbia Basin and Rocky Mountain Front areas and from 1 May - 20 June in the Intermountain area.

Incidental observations in unsuitable habitat such as ponds or large, low gradient (<1%) rivers not adjacent to known breeding sites, or observations on streams which have been identified as lacking breeding activity (e.g. migratory staging areas or stopovers).


## **SURVEYS AND BANDING**

### **MONTANA SURVEYS - 1998**

In 1998 ca. 54.1 km of streams were surveyed for Harlequin Ducks (Figure 1), in most cases multiple surveys of the same streams were conducted on different dates. Thus, a total of 235 stream km was surveyed in 1998 by MTNHP crews. Harlequin Duck pair surveys were conducted on 41.5 km of 6 streams by MTNHP crews: Rock Creek, Marten Creek, S. Fork Marten Creek, Swamp Creek, Vermilion River, Trail Creek (Flathead County). Glacier National Park personnel conducted a total of 81.4 stream km of pair surveys on 37.8 km of 9 streams (J. Ashley, pers. comm.). Brood surveys were conducted in July and August on 54.1 km of 7 streams (the above streams plus McDonald Creek, from Logan to Mineral creeks, in Glacier National Park) by MTNHP crews. Park personnel conducted brood surveys on 34.3 km of 3 streams.

**Lower Clark Fork** Two pair surveys were conducted between 9-22 May 1998 on each Lower Clark Fork stream (Rock Creek, Marten Creek/S. Fork Marten, Swamp Creek, Vermilion River). A minimum of 13 males and 11 females was observed on these four streams (Appendix 1). No adults were detected on Swamp Creek. Total numbers of ducks were about average on each of the major (four) streams except Swamp Creek. Swamp Creek has fluctuated between 0 and 3 females during 1992-1998. Light snow pack, early snow melt, and low stream flows aided our ability to do complete

**Figure 1. Streams surveyed for Harlequin Ducks in Montana in 1998, by the Montana Natural Heritage Program.**



pair surveys in May. These conditions probably had a strong influence on the length of time available to adults to locate suitable nest sites. We may have missed a few birds because they had earlier access to nesting sites than in average years.

Brood surveys were conducted between 4 July and 4 August 1998. Each of the four streams received 3-4 complete surveys. On the Lower Clark Fork streams, 3 females and 3 broods (13 young) were observed on Rock Creek. Based on the pair surveys of this stream, only 2 broods were anticipated, but the pair, lone male, and lone female seen on 19 May apparently represented adults of three pairs (see Appendix 1). Two females and 2 broods (11 young) were observed on Marten Creek, and 2 females and 2 broods (6 young during the final survey, 7 young on the previous survey) on the Vermilion River.

Previously marked birds identified in 1998 (see Appendix 2) in the Lower Clark Fork population included female blue VA on Rock Creek on 9 and 19 May (banded on 25 July 1995) and female blue TC on Rock Creek on 5 August (banded on Rock Creek on 26 July 1995). On Marten Creek, resighted birds on 10 May included male blue S1, male blue S8, male blue S9, female blue SE (banded on 26 April 1995, 10 May 1996, 10 May 1996, and 28 July 1995, respectively), female blue SA on 17 July (banded on 28 July 1995), and female blue SC on 4 August (banded 28 July 1995). On the Vermilion River, resighted birds included female blue X1 on 12 May (banded 29 July 1995), and female 925-09393 on 3 August (now also blue X2, banded on 31 July 1996).

**Glacier National Park.** Pair surveys were conducted during April and May 1998 by Glacier National Park personnel. A minimum of 16 females was observed on McDonald Creek, Glacier N. P. in 1998 (J. Ashley, pers. comm.).

During brood surveys, 5 females and 5 broods (24 young) were observed in the McDonald Creek drainage between 6 and 18 August (these include surveys conducted by Glacier N.P. personnel). One of these broods was encountered on Mineral Creek on 17 August, the remainder were on McDonald Creek proper. During the MTNHP brood survey of Upper McDonald Creek (Logan Creek to Mineral Creek) on 6 August, female blue G1 was captured with a brood, she had been banded on McDonald Creek the previous year (Appendix 2).

**Other Northwest Montana Areas.** A pair survey was conducted along 5 km of Trail Creek (N Fork Flathead River drainage, in Flathead County) on 14 May 1998. A minimum of 4 males and 3 females was observed. Confirmed resighted birds included female blue F7, originally banded on Trail Creek on 4 May 1996 (Appendix 2).

A brood survey was conducted along the same section of Trail Creek on 7 August 1998. One female with a brood of 1 young (Class IIC) was observed. The female (blue J8) was a bird originally banded on Trail Creek on 1 August 1996.

## SUMMARY OF MONTANA SURVEYS 1987-98

In Montana, over 5000 km of streams have been surveyed since 1987 (Reichel and Genter 1996, Reichel et al. 1997). Many of these stream reaches have been surveyed in multiple years and during both pair and brood seasons. Not all of these streams can be considered adequately surveyed. To be reasonably sure birds are not present on a stream where no previous sightings have occurred, at least two surveys should be conducted during 1-25 May. If more than one pair survey is done in a single year, they should be done at least 1 week apart. Lack of knowledge of proper survey timing resulted in many surveys prior to 1992 being done during June (after males had left and females are incubating) or

after 10 August when many birds in years of normal snow pack and runoff have left all streams but those in southwest Montana.

#### BANDING IN MONTANA: 1991-97

During 1998 in Montana, 2 adult males, 3 adult females, and 30 juveniles were captured and banded (Table 1); 1 male and 8 females were recaptured. This brings the total number of Harlequin Ducks banded since 1991 in Montana to 391 (61 males, 73 females, 257 juveniles).

**Table 1. Summary of Harlequin Ducks marked for the first time in 1998 (total ducks captured in all years including 1998 are in parentheses).**

Location	Male	Female	Juv.	Total
McDonald Creek drainage, Glacier NP*	(27)	1 (32)	19 (81)	20 (140)
Waterton River, Glacier NP		(1)	(7)	(8)
Trail Creek*	(10)	(9)	1 (18)	1(37)
Grave Creek		(3)	(6)	(9)
Spotted Bear River		(4)	(34)	(40)
Sullivan Creek, Flathead Co.		(2)	(10)	(12)
Cache Creek	(1)			(1)
Blackfoot River, North Fork		(1)	(2)	(3)
Marten Creek, Sanders Co.*	2 (17)	1 (7)	5 (43)	8 (67)
Rock Creek, Sanders Co.*	(4)	(5)	2 (19)	2 (28)
Swamp Creek, Sanders Co.*	(1)	(2)	(12)	(15)
Vermilion River, Sanders Co.*	(1)	1 (5)	3 (25)	4 (31)
<b>TOTAL</b>	<b>2 (61)</b>	<b>3 (73)</b>	<b>30 (257)</b>	<b>35 (391)</b>

\* streams surveyed in 1998.

## MOVEMENT

#### ON THE BREEDING GROUNDS

In Montana and Idaho, several relatively long-distance movements have been documented both within and between years (Table 2). One inter-stream movement was documented in 1998, in the Lower Clark Fork drainage. A female banded as a juvenile on Swamp Creek on 27 July 1995 was observed with an unmarked adult male on 9 and 19 May 1998 on Rock Creek, a movement of about 21 km (along water courses, not a direct line). We have recorded movements between Marten Creek and the three other streams regularly surveyed in the Lower Clark Fork drainage (Rock Creek, Swamp Creek, Vermilion River), as well as with West Gold Creek in northern Idaho. The Swamp Creek to Rock Creek movement was the first between streams other than Marten Creek in this population. Marten Creek appears to be an extremely significant area for Harlequin Ducks of the Lower Clark Fork population, but some level of movement between all of these streams is likely, as indicated by the movement observed in 1998.

**Table 2. Significant movements of Harlequin Ducks within and between years on the breeding grounds in Montana and Idaho (Cassirer and Groves 1994, Reichel and Genter 1994, 1995; Ashley 1995, 1996; Reichel et al. 1997; Hendricks and Reichel 1998; Cassirer pers. comm.).**

Age and sex when banded	1st Date	Location	2nd Date	Location	Km moved
Adult Male	1990	Gold Creek, ID	1991	Granite Creek, ID	14
Adult Male 755-76075	5/26/93	Marten Creek, Devils Gap	4/27/95	Vermilion River, 0.1 mi above Miners Gulch	31
Juv. Female 805-90262 changed to 925-09364	8/10/92	West Gold Creek at Lake Pend Oreille, ID	7/29/96	Marten Creek, near mouth	50
Adult Female 755-76007	8/4/92	Marten Creek, mouth of (w/ brood)	7/30/93	Swamp Creek, T25N R31W Section 9 (w/ brood)	16
Adult Female 755-76025	8/10/92	McDonald Creek above McDonald Lake (w/ brood), Glacier N P	6/29/95	Middle Fork Flathead River (w/ brood)	18
Adult Female 755-76013	7/28/95	Marten Creek, near mouth (with 6 young 925-09336, 37, 38, 39, 40, 41)	7/29/95	Vermilion River, near Sims Creek confluence (with same 6 young) T	26
Adult Female 925-09374	7/31/96	Grave Creek, 0.7 mi above Cat Creek, Lincoln Co	8/20/96	McDonald Creek, near McDonald Falls, Glacier National Park	75
Juv. Female 775-37376	7/26/95	Rock Creek, T26N R32W S22SE	8/9/97	Marten Creek, near Devils Gap	21.5
Juv. Female 775-37379	7/25/95	Swamp Creek, T26N R31W S34SESWSW	5/9/98 and 5/19/98	Rock Creek, T26N R32W S27NWNE (paired, probably raised a brood)	21

The two longest movements to date were recorded in 1996. A female marked as a juvenile on West Gold Creek in Idaho in 1992 was found injured and without a brood on Marten Creek in 29 July 1996 (a 50 km movement). There have been no documented cases of females breeding on streams farther than 20 km from their natal stream (always within the same drainage), although the 1998 observation indicates breeding probably occurred on a stream other than the natal stream in the Lower Clark Fork population. The second long distance movement was a female marked on Grave Creek on 31 July 1996 without a brood and observed again, 75 km away, on McDonald Creek on 20, 22, and 28

August 1996 (Ashley 1997). This was likely a post-breeding exploratory movement or wandering during migration. Since no surveys had been done on Grave Creek since May of that year, it is unknown if she spent the summer there.

Both sexes of Harlequin Duck have been observed using different nearby drainages during different years in Montana and adjacent Idaho (Table 2). Two cases of within-year drainage change by adult females have also been noted. These observations indicate that movements of up to 30 km within an aggregate of streams may occur routinely but infrequently, both within and between years. Movements occurred over large reservoirs (Noxon Reservoir) and lakes (Lake McDonald). The 1995 movement by a female and her entire fledged brood to the Vermilion River was likely the result of disturbance due to marking. However, the movement took place at least 4 hours following the release of the birds. A breeding female (755-76025) in Glacier Park has been seen at several locations on different streams in the park over the 4 years since her banding (Ashley 1995); locations in Table 2 are the maximum distance moved during the 4 year period.

There is little published literature regarding movement within the breeding grounds. Kuchel (1977) found that pairs used lower McDonald Creek prior to establishing home ranges higher up along the stream. Once established, pairs rarely moved more than 1-2 km, although movements of up to 8 km were recorded. Kuchel (1977) found unpaired males moved considerably more during the breeding season, with movements of up to 10 km recorded. In a reanalysis of Kuchel's (1977) data, Cassirer and Groves (1992) found that linear home ranges during the breeding season averaged 7.7 km ( $SD = 2.34$ ) on McDonald Creek, similar to the 7 km reaches used in Idaho.

On the Bow River in Banff National Park, 5 pairs of birds were marked at what is probably a staging area or local migratory corridor (Smith 1996). Two pairs remained in a 2 km section of river where they were banded, and another remained in a 2 km stretch about 12 km downstream; one pair remained within about 6 km until the female moved about 8 km up a drainage, perhaps to breed; the final pair moved about 15 km downstream within 22 days (Smith 1996).

For 35 Harlequins marked in Iceland, Bengtson (1972) found no movement between breeding streams and movement of only a few km within drainages. Not only did the birds return to the same drainage, but in 22 out of 33 cases, the birds were observed within 100 m of their locations during the previous year (Bengtson 1972).

## MIGRATION

*Nature of migration.* All inland populations of Harlequin Duck migrate to coastal waters. A marked female seen on Granite Creek, Idaho on 17 July 1991 was relocated 13 days later near Battleship Island in the San Juan Islands, Washington (Cassirer and Groves 1992). In Iceland, birds are thought to swim up the rivers from the coastal wintering grounds to the freshwater breeding sites (Gudmundsson 1961 *in* Bengtson 1966).

Sibling juveniles may migrate together to the coast, as indicated by the presence of 3 siblings at Hornby Island, B.C. (Harlequin Duck Working Group Database, G. Schirato, Wash. Dept. Fish & Wildlife pers. comm.) which were marked together 7 months earlier on Swamp Creek, Montana. Whether females and their broods migrate together in some instances is unknown. However, it is known that females occasionally leave prior to their young fledging. In Montana, out of 135 broods observed during 1988-98, 16 broods (11.9%) were found without the hen prior to migration (Reichel et al. 1997, Hendricks and Reichel 1998, Ashley pers. comm.). Age class of the 16 abandoned broods when they were first observed alone was as follows: 2 were Class I, 3 were Class II, 3 were Class III, and 8 broods were first observed without the adult female following fledging. In one additional case, a

brood of 7 was marked with the female on 11 Aug 1992, on 2 September the female was seen with 5 of her fledged juveniles, while one of the brood was observed alone 2.5 km away. Adult males apparently return to coastal sites shortly after breeding (June-July), where they complete prebasic molt by August, and then undergo pre-alternate shortly thereafter (Cooke et al. 1997), attaining alternate plumage by October. Pair formation begins at this time and extends well into winter (Robertson et al. 1998a). Although timing of molt was not related to pairing success, males that molt quickly into the bright alternate plumage have a higher probability of pairing with females in the winter and spring coastal aggregations (Robertson et al. 1998b).

**Coastal wintering sites.** Of 249 Harlequins banded in Montana from 1991-1995, a minimum of 24 have been reported from Oregon (2), Washington (1), and southern British Columbia (21) through September 1996 (G. Shirato pers. comm.). Sexes and ages at banding show the following numbers and percentages observed: adult females (6, 11%), adult males (2, 5%), juvenile females (9, 7%), and juvenile males (7, 5%).

Birds banded during 1996-1998 are not included in the above summary of resighting records for birds marked in Montana. Nevertheless, during 1997 to January 1999 more than 30 individuals banded in Montana have been reported from British Columbia (Vancouver and Hornby islands) and at least 2 birds from Washington (Harlequin Duck Working Group pers. comm., numerous observers). Resightings include 7 adult males, 5 adult females and 2 juveniles from the Lower Clark Fork streams: Rock Creek (2 males, 1 female, 2 juveniles), Swamp Creek (3 males, 2 females), Marten Creek (1 male, 1 female), Vermilion River (1 male, 1 female). Additional coastal resightings during 1997-1998 for birds banded on other Montana streams included McDonald Creek (7 males, 3 females), Trail Creek (1 male), Spotted Bear River (1 female) and North Fork of the Blackfoot River (1 male, 1 female). Female VA, the Swamp Creek bird seen paired on Rock Creek in May 1998 (Table 2) and probably producing a brood on Rock Creek, was reported off Vancouver Island, British Columbia on 11 September 1998 (Ken Wright pers. comm.).

**Migratory behavior.** It is believed that nearly all one-year-old birds, and some (perhaps most) two-year-old birds remain in coastal water, not moving to breeding streams until they are 2-4 years of age. The proportion of each age class that stays on the coast has yet to be determined, but indications are that one-half of 2-year-old females and one-quarter of 3-year-old females do not return to the breeding grounds. Five 2-year-old females were seen during 1997 surveys of the Lower Clark Fork streams. Four of these females were on their natal stream (Marten Creek) and 3 were siblings. The fifth female was born on Rock Creek and was encountered on Marten Creek (see Table 2). No two-year-old males or females were noted during the 1998 surveys. Wallen (1987) reported that a 1-year-old female (of 11 total) returned to her natal stream (Upper Moose Creek) in Grand Teton National Park in 1986. This is the only report of a 1-year-old female on the breeding grounds. No one- or two-year-old males, out of 246 observations of males, have been seen in Montana during 1992-96 surveys (Table 7, Ashley pers. comm.).

## **DEMOGRAPHY AND POPULATIONS**

### **MEASURES OF BREEDING ACTIVITY**

**Age at first breeding; intervals between breeding.** Only a single known-aged male has been seen with a mate; it was marked by MTNHP as a juvenile in 1992 on Mineral Creek (Glacier National Park), Montana, and observed by J. Ashley paired with a female (white NH) at Hornby Island, BC, in

March 1996. Adult male breeding plumage is attained at three years of age (Phillips 1925). No one- or two-year-old males, out of 246 independent male observations, have been observed in Montana during 1992-96 surveys (Ashley pers. comm.).

The youngest female known to have bred is a 2-year-old that raised a brood of 3 in 1994 on Trail Creek, Montana. Several 3-year-old females have bred successfully in Montana, including 4 in 1998 (TC on Rock Creek and probably VA; SA and SC on Marten Creek). Only a single 4-year-old is known to have bred successfully, through 1997 (on Marten Creek in 1996). One 5-year-old female is known to have bred successfully on her natal stream (McDonald Creek in 1997).

Some females on breeding streams apparently do not lay eggs (Bengtson and Ulfstrand 1971, Dzinbal 1982, Wallen 1987, Cassirer and Groves 1991). Bengtson and Ulfstrand (1971) examined ovaries of 6 non-breeding females and reported that none had laid eggs. They reported that 15-30% ( $n=48$ ) of adult females (based on bursae inspection) were non-breeders. Many of these non-breeding "adults" may have been young (2-3 year-old) birds, since cloacal examination gives adult status to 2-year-olds.

**Annual and lifetime reproductive success.** Reproductive success in Montana was above average in 1998, with one of the larger number of broods and young per female recorded, especially on many Lower Clark Fork streams and McDonald Creek (Table 3, 4). Values for streams surveyed in 1998 (0.41 broods per female, 1.72 young per female, 4.23 young per brood) were similar to values for 1992 (Table 4: 0.55 broods per female, 1.37 young per female, 3.38 young per brood), although young per brood was the second largest reported.

The data for 1998 and other years (Table 4) are not directly comparable, however, because not all streams represented in the pooled annual survey data were identical. Streams with the largest duck populations have the greatest influence on mean values, and there is no evidence that production on any single stream or subset of streams is typical each year for all breeding streams in the state. Pooling results for the entire state, as presented in Table 4, also masks fluctuations for particular streams or drainages. This is especially evident for the Lower Clark Fork streams (Appendix 3). Each of these streams has experienced at least one year during 1992-1998 where no broods were produced, but in only one year (1992) did the majority of streams fail to produce broods. Interestingly, during that year Marten Creek produced more broods (4) than in any other year for which there are data.

Until data are available on age-specific reproduction and longevity, no lifetime reproductive success can be calculated. Therefore, this has not been calculated for any Montana population.

**Proportion of females that rear broods to nest-leaving.** Harlequin Ducks raise only a single brood each year. The proportion of females successfully raising a brood in a single year varies widely between years (Table 4). In Montana during 1998, 41% of 32 females successfully raised a brood. Stream surveys between 1974 and 1998 found that 409 females raised 151 broods for a total 37% of females with broods (range 7-55%), and an annual mean of 35% for 12 years of data. Throughout their range, the percentage of females which successfully raise broods varies from 7-56% (Bengtson and Ulfstrand 1971, Kuchel 1977, Wallen 1987, Cassirer and Groves 1991, Reichel et al. 1997).

**Sex ratio.** In 1998 on the Lower Clark Fork streams, we counted 13 males and 11 females during May pair surveys. These numbers are probably low by 2 males and 2 females. Three broods were produced on Rock Creek (Appendix 1) even though we counted only enough birds for 2 pairs, and a marked female with a brood on the Vermilion River was not counted during pair surveys. Therefore, sex ratio on these streams was 1.15:1 (m:f) in 1998. During the spring pair season for all years combined through 1997, a sex ratio of 1.52:1 has been observed in Montana (m:f,  $n=786$ ; 474 males:312 females). These values are based on independent male observations during the period 27



**Table 3. Harlequin Duck reproduction in 1998 for Montana streams with both pair and brood (near fledging) surveys.**

Stream	#Adult Females	#Broods	#Young
<b>Flathead Drainage</b>			
McDonald Creek <sup>a</sup>	16	5	24
Trail Creek	3	1	1
Drainage Total	19	6	25
0.32 Broods per adult female			
1.32 Young per adult female			
4.17 Young per brood			
=====			
<b>Lower Clark Fork Drainage</b>			
Marten Creek	7	2	11
Rock Creek	3	3	13
Swamp Creek	0	0	0
Vermilion River	3	2	6
Drainage Total	13	7	30
0.54 Broods per adult female			
2.31 Young per adult female			
4.28 Young per brood			
=====			
<b>GRAND TOTAL</b>	<b>32</b>	<b>13</b>	<b>55</b>
0.41 Broods per adult female			
1.72 Young per adult female			
4.23 Young per brood			

<sup>a</sup> from Ashley (pers. comm.)

April - 31 May and the maximum number of females seen when more than one survey was done during a single season on a single stream (see Reichel et al. 1997 for more details). Cassirer (1995) found a spring adult sex ratio of 1.31:1 (m:f, n = 81) in 1995 on Idaho streams. In Banff National Park, Alberta, sex ratios varied from 1.37:1 in May to 1.81 in June (Smith 1996). In Iceland, sex ratios on the breeding grounds varied from 1.17 - 2.33:1 during 5 summers in late May - early June (Bengtson 1966, Bengtson 1972, Gardarsson 1979).

In coastal British Columbia, the apparent sex ratio is 1.5:1 (544 birds) in winter, declining from 1.4:1 (297 birds) in March-April (Campbell *et al.* 1990), this grows to 4.3:1 in May, and by July, when adult females are still on the breeding streams, it reaches 18.2:1 (n = 1633 birds).

**Table 4. Harlequin Duck reproductive parameters 1974-75 (Kuchel 1977) and 1989-1998 for streams with both pair and brood surveys.**

Year	No. adult females	No. broods	No. young	broods per ad. female	young per ad. Female	young per brood
1974	11	3	12	0.27	1.09	4.00
1975	15	1	2	0.07	0.13	2.00
1989	13	7	41	0.54	3.15	5.86
1990*	31	17	65	0.55	2.10	3.82
1991*	37	9	31	0.24	0.84	3.44
1992*#	71	39	132	0.55	1.37	3.38
1993#	49	21	59	0.43	1.20	2.81
1994#	30	10	40	0.33	1.33	4.00
1995#	48	11	42	0.23	0.87	3.82
1996*#	44	14	56	0.32	1.27	4.00
1997 <sup>a</sup>	28	6	25	0.21	0.89	4.17
1998 <sup>a</sup>	32	13	55	0.41	1.72	4.23
Total	409	151	560	0.37	1.37	3.71

Annual mean±SD

0.35±0.15    1.33±0.75    3.79±0.92

\* includes data from the Rocky Mountain Front (Diamond and Finnegan 1992, 1993; D. Whittekind, pers. comm.)

# includes data from Ashley (1994a, 1994b, 1995, 1996, 1997, pers. comm.)

## FIDELITY TO BREEDING STREAMS

**Adult fidelity to breeding stream.** In Montana, 391 Harlequins (61 adult males, 73 adult females, 257 juveniles) have been banded from 1991 through 1998 (Table 1). In 1998, 3 males and 12 females originally marked as adults were resighted on the streams on which they had been banded in a previous year (Appendix 2). This figure includes females in Glacier National Park, but not males. Streams included Marten Creek (3 males), Vermilion River (2 females), Trail Creek (2 females), and McDonald Creek (8 females). Through 1996, 60% of males (25 of 42) and 64% of females (36 of 56) returned at least 1 year following marking. Figures have not yet been calculated using 1997-1998 data. Lower return rate of males may be a result of mating with new females on the coast and following them to female natal streams.

**Fidelity to natal stream.** In 1998, three 3-year-old females returned to natal streams (Rock Creek and Marten Creek) and successfully raised broods. One 3-year-old female returned to a stream (Rock Creek) not her natal stream (Swamp Creek) and probably raised a brood.

## CONSERVATION AND MANAGEMENT

### IMPLICATIONS OF THE CURRENT RESEARCH

This section essentially restates what has already been presented in previous reports (e.g., Reichel et al 1997, Hendricks and Reichel 1998). These comments are worth presenting again to emphasize that many populations continue to be exposed to serious threats. Much of what is known

about the life history highlight the precarious nature of the populations in the Rocky Mountains of the United States. Factors that limit recolonization and increase the possibility of extirpation include: 1) high female natal site fidelity; 2) high adult site fidelity; 3) pair bonds developing on the wintering grounds; 4) low levels of movement on the breeding grounds; 5) relatively advanced age at first reproduction; 6) little chance of renesting after about 2 weeks following the start of incubation; 7) low and irregular levels of reproductive success; 8) patches of suitable habitat which are highly fragmented; 9) sensitivity to disturbance; 10) the clumped distribution of pairs, even in apparently homogeneous habitat; 10) declining range-wide and regional population levels; 11) relatively small and isolated regional populations; and 12) use of coastal wintering habitat immediately offshore (often less than 100 m).

Harlequin Ducks apparently form pair bonds on the coast. Females apparently lead their mates to the breeding streams. Site fidelity is high, both for first-time and experienced breeders, and probably exceeds 90% for both categories. This leaves very few birds to explore and "pioneer" new sites. Harlequin Ducks may key in on areas with other ducks present, like many other bird species with clumped distributions. In other words, good habitat to a "pioneer" is where ducks are already present; empty habitat would be unlikely to be colonized. Suitable habitat in the Rocky Mountains is currently sparse and widely separated. Much has probably been lost and fragmented by alteration of stream-side land and building of reservoirs.

Small breeding populations face several challenges. Random events, such as several birds dying or several poor reproductive years caused by flooding, can dramatically reduce already small populations or eliminate them. Females do not breed until 2 or more years-old and adult success rates may not occur until 4 or more years-old. This means that mortality must be low or few ducks will even make it to breeding age. Once a duck attains breeding age, it can only produce a single brood each year. While many bird species renest when the nests or young are lost, there is little possibility of Harlequins renesting after more than a week or two following egg laying. Males return to the coast a week or two after the females begin incubation. The result of the above factors is that reproductive success is low and highly variable. An average female is probably 4 or 5 years old before she has raised even 2 female ducklings to fledging. It is likely that mortality in the first 6 months following fledging is high.

Harlequins from the U.S. Rocky Mountains move to the Pacific coast off Oregon, Washington, and British Columbia following breeding and remain there until the following spring. They are concentrated in areas with rocky shorelines. Harlequins are the most coastal of wintering sea ducks and are thus more susceptible to hunting and oil spills than most sea ducks.

This set of facts does not bode well for the Harlequin Duck in Montana, where 10 of 27 occurrences consist of only 1 or 2 pairs of ducks. It also shows the critical nature of the 6 occurrences with more than 15 pairs as a source of stability to the Rocky Mountains regional population.

## **PRIORITIES FOR FUTURE RESEARCH**

The following are among the top future research priorities and are primarily a subset of those listed by the Harlequin Duck Working Group (1993) and by Cassirer *et al.* (1996).

## 1) What are the impacts of human disturbance on breeding and wintering Harlequin Ducks?

Several independent studies have documented the sensitivity of Harlequin Ducks to human disturbance, primarily through the relationship of sighting locations to the accessibility of those locations (Kuchel 1977, Wallen 1987, Diamond and Finnegan 1993, Cassirer and Groves 1991, 1994, Clarkson 1992, Ashley 1994). Specifically, boating has been shown to have a significant negative correlation with numbers of ducks present in one area on a medium-sized stream (Clarkson 1992, Hunt 1993). Observations in other areas tend to support this conclusion (Cassirer and Groves 1991, Brady pers. comm. *in* Clarkson 1992) though it may not be the case in very large streams (Smith 1996, 1999). Fishing and human presence have also been suggested as causes of disturbance. However, although specific examples exist for both disturbances, statistical data analyses are lacking (Wallen 1987, McEneaney 1994, Cassirer and Groves 1991).

Other than for boating (Clarkson 1992, Hunt 1993), wide-scale analyses have not yet been attempted nor have analyses of the effects of most specific kinds and amounts of human activities. Several specific studies should be performed to address these questions.

Initially, wide-scale data on Harlequin streams are required. Data needed on the ducks themselves include productivity and population size on each stream. Data needed on the streams used by Harlequin Ducks include length of stream segments used during pair and brood seasons, categories and locations of land ownership, hydrogeological properties, habitat characteristics, and current human use (by roads, trails, structures, activity, etc.). A first step will be to see which of this information is already available and which is lacking that needs gathering in the field. Unused and/or unknown streams that fit physical parameters of used streams can then be selected and compared in respect to kind and amounts of disturbance/accessibility.

Following wide-scale analyses, Harlequin response to humans requires evaluation. Initial responses to surveyors could be recorded. Note that this would only provide immediate, in-sight response of birds seen. Presumably some birds would react prior to the surveyor seeing them and thus not be observed at all. Nor would such a study reveal length of time or distance moved in reaction to disturbance. A more precise but intrusive method would be to use radio-telemetry on the birds. Radio-telemetry would also provide more accurate data on use of habitat types and locations relative to human development/access points.

Finally, when actions are taken on Harlequin streams, monitoring to determine effects of those actions should be implemented, thereby providing for adaptive management and prevention of future mistakes. Specific land management or development actions on Harlequin streams should be preceded by at least two years of baseline marking and surveying for population size and productivity, areas used at different seasons, habitat evaluation, and pre-action levels of human activity and development. Monitoring should continue to occur during and following the action. Actions needing special attention include road, campsite, and trail construction activity, including any increased accessibility and changes in human use of the area. Actions that might result in changes to flow regimes or water quality include mining, road building, timber harvest, industrial development, and water/hydroelectric development. Also requiring attention are changes in fishing regulations, which could alter fisherman impact on an area, and building of structures, such as industrial areas, dams, or houses, which will increase access to and use of a Harlequin stream. Possibilities for mitigation and habitat restoration can be explored during these projects.

## 2) What is the extent and nature of movements in breeding and wintering areas?

This information is needed to determine the possibilities for natural recolonization of new and historic Harlequin occurrences, natural supplementation of existing occurrences, particularly small populations, and the strength of natal and adult fidelity to particular sites. This information is necessary for successful modeling of Harlequin populations and their stability, using both breeding and wintering ground data.

Radio-telemetry may give quick results from the standpoint of local daily movements (see Smith 1999), however, long distance (>5 km) movements may be relatively rare, and with limited numbers of ducks radioed, may not be best for long distance movement detection. For long distance and moves between years, visibly marking birds is best.

Determining fidelity to natal areas will be a long-term project. Montana has the strongest start, with 391 birds banded on the breeding grounds since 1992. Data so far indicate high fidelity to natal streams.

Much data is now available in relation to wintering ground movements and fidelity, and additional data is currently being collected in Washington, Alaska, and British Columbia. Sufficient information for use in detailed population modeling is currently being processed (Robertson pers. comm.). For an accurate model, data are necessary from both the breeding and wintering grounds.

## 3) Are distinct metapopulations (such as a Rocky Mountain breeding population) identifiable within the Pacific range of the Harlequin Duck?

Knowledge of the degree of genetic differences among and within wintering and breeding subpopulations would allow an assessment of the appropriate management units for various Harlequin conservation strategies. Dan Esler, Alaska National Biological Service, and Maggie Brown (Department of Wildlife, Fish and Conservation Biology, University of California - Davis) are currently examining this question (see Brown 1998).

## 4) What are the critical habitat components limiting Harlequin Duck breeding and wintering populations?

Harlequin Ducks use a wide variety of habitats on the breeding grounds, from forests to tundra. Habitat usage should be documented over a large number of study areas to identify common habitat components for comparison to available habitat, both large and small scale components should be considered.

## 5) How and why do productivity and survival change over time and different areas, and what are the relative impacts of these changes on populations?

Long term studies are needed to determine population parameters for incorporation into population models (with information from movements on the breeding and wintering grounds). Needed population parameters include: productivity; age-related survival, recruitment, age(s) at first breeding and/or successful breeding, age(s) last breeding, life expectancy; and causes and timing of mortality. This information will be acquired only through long-term studies involving marked birds on both the breeding and wintering areas. We are currently in an optimum position to complete

studies needed on the breeding grounds, with 7 years of data on portions of the Montana breeding population. Combined with the continued marking and study of coastal populations by Alaska, Washington, Oregon, and British Columbia, and interior populations in Alberta, many of these parameters may be known in the next few years.

The most difficult question to be answered involves the causes of mortality, which is not tractable given current technology. If and when small, long range mortality transmitters are available for ducks, this topic should be pursued. Some females are killed on nests (Smith 1999) by predators, but the significance of predation is not yet clear.

6) What are the characteristics of Harlequin Duck migration? How well defined are migratory staging areas and migration corridors?

This question may not be tractable given current technology. When small, long range mortality transmitters are available for ducks, this topic should be pursued. Some answers may come from large scale marking of individuals, and perhaps by relocating radioed birds. We still have few reports of Harlequin Ducks moving between the coast and inland breeding areas. Nor do we know routes of movement on a smaller landscape scale. For example, Harlequin Ducks that breed on the east slope of the Continental Divide must cross the Divide on their movements to and from the coast, but where do they do so? Do certain drainages serve as "funnels" for most ducks in local populations during migratory movements?

## **REFERENCES**

- Alford, C. E. 1920. Some notes on the harlequin duck. *British Birds* 14:14-17.
- American Ornithologists' Union. 1983. Check-list of North American Birds, 6th edition. Allen Press, Inc., Lawrence, Kansas. 877 pp.
- Ashley, J. 1992. A summary of documented harlequin duck observations in Glacier National Park, 1874-1992. Unpublished draft report, 18 pp. plus maps.
- Ashley, J. 1994a. 1992-93 harlequin duck monitoring and inventory in Glacier National Park, Montana. Unpublished report. Div. Res. Manage., Glacier Natl. Park, Montana. 57 pp.
- Ashley, J. 1994b. Progress report: harlequin duck inventory and monitoring in Glacier National Park, Montana. Unpublished report. Div. Res. Manage., Glacier Natl. Park, Montana. 14 pp.
- Ashley, J. 1994c. Status of Harlequin ducks in Glacier National Park, Montana. P. 2 in: Proc. 2nd Ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Ashley, J. 1995. Harlequin duck surveys and tracking in Glacier National Park, Montana. Unpublished report. Div. Natural Resources, Glacier National Park, West Glacier, Montana. 41 pp.
- Ashley, J. 1996. Harlequin duck inventory and monitoring in Glacier National Park, Montana. Unpublished report. Div. Natural Resources, Glacier National Park, West Glacier, Montana. 21 pp.
- Ashley, J. 1997. Harlequin Duck inventory and monitoring in Glacier National Park, Montana. Unpublished report. Division of Natural Resources, Glacier National Park. West Glacier, Montana. 36 pp.

- Atkinson, E. C. and M. L. Atkinson. 1990. Distribution and status of harlequin ducks (*Histrionicus histrionicus*) on the Targhee National Forest. Idaho Dept. of Fish and Game, Nongame and Endangered Wildl. Prog. 25 pp.
- Bailey, A. M. and R. J. Niedrach. 1965. Birds of Colorado. Vol. 1. Denver Mus. Nat. Hist. 454 pp.
- Bailey, A. M. and R. J. Niedrach. 1967. A pictorial checklist of Colorado birds, with brief notes on the status of each species in neighboring states of Nebraska, Kansas, New Mexico, Utah, and Wyoming. Museum of Natural History, Denver.
- Bailey, V. 1918. Wild animals of Glacier National Park: the mammals, with notes on physiography and life zones. U.S. Department of the Interior, National Park Service. 210 pp.
- Baldwin, S. P., H. C. Oberholser, and L. G. Worley. 1931. Measurements of birds. Sci. Publ. Cleveland Mus. Nat. Hist. Vol. 2. 165 pp.
- Bauer, K. M. and U. N. Glutz von Blotzheim. 1969. Handbuch der Vogel Mitteleuropas. Band 3. Frankfurt am Main. Akademische Verlagsgesellschaft.
- Bellrose, F. C. 1980. Ducks, geese and swans of North America. Stackpole Books, Harrisburg, Pennsylvania. 540 pp.
- Bengtson, S.-A. 1966. Field studies on the harlequin duck in Iceland. Wildfowl Trust Ann. Rep. 17: 79-84.
- Bengtson, S.-A. 1972. Breeding ecology of the harlequin duck (*Histrionicus histrionicus*) in Iceland. Ornis Scand. 3: 1-19.
- Bengtson, S.-A. and S. Ulfstrand. 1971. Food resources and breeding frequency of the harlequin duck *Histrionicus histrionicus* in Iceland. Oikos 22: 235-239.
- Bent, A. C. 1925. Life histories of North American wild fowl. Order: Anseres (Part II). U.S. Natl. Mus. Bull. 130. Washington, D.C. 316 pp.
- Boertmann, D. 1994. An annotated checklist to the birds of Greenland. Meddr. Grønland, Biosci. 38: 63 pp.
- Brazil, M. A. 1991. The birds of Japan. Smithsonian Institution Press, Washington, D.C. 466 pp.
- Breault, A. M. 1993. Harlequin duck status report 1992. British Columbia. Pp. 60-64 in: Cassirer, E. F., et al., (eds.), Status of Harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Breault, A. M. and J.-P. L. Savard. 1991. Status report on the distribution and ecology of harlequin ducks in British Columbia. Can. Wildl. Serv., Pac. and Yukon Reg., Tech. Rep. Series 110. 108 pp.
- Breault, A. M. and K. M. Cheng. 1990. Use of submerged mist nets to capture diving birds. J. Field Ornithology. 6: 328-330.
- Brown, M. E. 1998. Population genetic structure, kinship, and social associations in three Harlequin Duck (*Histrionicus histrionicus*) breeding subpopulations. M.S. Thesis, University of California, Davis. 104 pp.
- Burleigh, T. D. 1951. Spring migration. Audubon Field Notes 5: 266-268.
- Burleigh, T. D. 1952. Spring migration. Audubon Field Notes 6: 258-260, 291, 292.
- Burleigh, T. D. 1972. Birds of Idaho. The Caxton Printers, Ltd., Caldwell, ID. 467 pp.
- Bustnes, J. O. and K. E. Enkstad. 1990. Effects of patagial tags on laying date and egg size in common eiders. J. Wildl. Manage. 54(2): 216-218.

- Byrd, G. V., J. C. Williams, and A. Durand. 1992. The status of Harlequin ducks in the Aleutian Islands, Alaska. Pp. 14-22 in: Proc. Harlequin duck Symposium., April 23-24, 1992, Moscow, Idaho. ID Dept. of Fish & Game. U.S. For. Serv. Intermountain. Res. Stat., ID Panhandle Nat. Forests, and NW Sect. of Wildl. Soc. 46 pp.
- Cahn, A. R. 1947. Notes on the birds of the Dutch Harbor area of the Aleutian Islands. Condor 49:78-82.
- Campbell, R. W., N. K. Dawe, I. McTaggart-Cowan, J. M. Cooper, G. W. Kaiser and M. C. McNall. 1990. The birds of British Columbia, Vols. 1 and 2: Non passerines. Royal British Columbia Museum, Victoria, B. C. 518 and 636 pp.
- Carlson, J. C. 1990. Results of harlequin duck (*Histrionicus histrionicus*) surveys in 1990 on the Flathead National Forest, Montana. Unpublished report. 31 pp.
- Cassirer, E. F. 1989. Distribution and status of Harlequin ducks (*Histrionicus histrionicus*) on the Nez Perce National Forest, Idaho. Report on Challenge Cost Share Project. 13 pp.
- Cassirer, E. F. 1993. Harlequin duck status report 1992: Idaho. Pp. 27-30 in: Cassirer, E. F., et al., (eds.), Status of Harlequin ducks in North America. 83 pp.
- Cassirer, E. F. 1994. Proposed inventory and monitoring protocol for harlequin ducks in northern Idaho. Paper presented at Interagency Rare Animal Workshop, March 2, 1994, Post Falls, Idaho. 14 pp.
- Cassirer, E. F. 1995. Harlequin duck monitoring on the Moyie River and other tributaries to the Kootenai River in northern Idaho subsequent to natural gas pipeline construction. Unpublished report. Idaho Department of Fish and Game, Lewiston, ID. 11 pp.
- Cassirer, E. F. and C. R. Groves. 1989. Breeding ecology of harlequin ducks (*Histrionicus histrionicus*) on the Kaniksu National Forest, Idaho. Report on Challenge Cost Share Project. 48 pp.
- Cassirer, E. F. and C. R. Groves. 1990. A summary of harlequin duck sightings in Idaho, 1989. Unpubl. rep. Idaho Dep. of Fish and Game, Boise. 11 pp.
- Cassirer, E. F. and C. R. Groves. 1990. Distribution, habitat use and status of harlequin ducks (*Histrionicus histrionicus*) in northern Idaho, 1990. Idaho Dept. of Fish and Game, Nongame and Endangered Wildl. 54 pp.
- Cassirer, E. F. and C. R. Groves. 1991. Harlequin duck ecology in Idaho: 1987-1990. Idaho Dep. of Fish and Game, Boise. 93 pp.
- Cassirer, E. F. and C. R. Groves. 1992. Ecology of Harlequin Ducks in northern Idaho; progress report 1991. Idaho Dept. Fish and Game, Boise, ID. 74 pp.
- Cassirer, E. F. and C. R. Groves. 1994. Breeding ecology of Harlequin ducks in Idaho. P. 3 in: Proc. 2nd Ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Cassirer, E. F. and C. R. Groves. 1994. Ecology of harlequin ducks (*Histrionicus histrionicus*) in Northern Idaho. Study No. 4202-1-7-2. Idaho Dept. of Fish and Game. 51 pp.
- Cassirer, E. F. and G. Schirato. 1990. Harlequin duck boat surveys, northwest Washington coast, 9/24-9/29/90. Washington Dept. of Wildl. 1 p.
- Cassirer, E. F., G. Schirato, F. Sharpe, C. R. Groves, and R. N. Anderson. 1993. Cavity nesting by harlequin ducks in the Pacific Northwest. Wilson Bull. 105:691-694.
- Cassirer, E. F., J. D. Reichel, R. L. Wallen, and E. Atkinson. 1996. Harlequin Duck (*Histrionicus histrionicus*)



conservation assessment and strategy for the U.S. Rocky Mountains Draft report

- Cassirer, F. 1995. Harlequin duck monitoring in northern Idaho, 1995. Cooperative project report. Idaho Dept. of Fish & Game, North Idaho Traditional Bowhunters, U.S. Forest Service, and Washington Department of Fish and Wildlife. 20 pp.
- Chadwick, D. H. 1992. Some observations of a concentration of harlequin ducks in the Strait of Georgia, British Columbia. Pp. 33-40 in: Proceedings Harlequin Duck Symposium, Apr 23-24, 1992, Moscow, ID. 45 pp.
- Childs, H. E., Jr. 1969. Birds and mammals of the Pitmegea River region, Cape Sabine, northwestern Alaska. University of Alaska Biological Papers, No. 10. 74 pp.
- Clarkson, P. 1992. A preliminary investigation into the status and distribution of harlequin ducks in Jasper National Park. Unpubl. Tech. Rep. Nat. Res. Conserv., Jasper Natl. Park, Alberta. 63 pp.
- Clarkson, P. 1994. Managing watersheds for harlequin ducks. Unpublished presentation. American River Management Society, River Without Boundaries Symposium, Grand Junction, CO. 33 pp.
- Clarkson, P. and R. I. Goudie. 1994. Capture techniques and 1993 banding results for moulting Harlequin ducks in the Strait of Georgia, B.C. Pp. 11-14 in: Proc. 2nd Ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Cooke, F., G. J. Robertson, R. I. Goudie and W. S. Boyd. 1997. Molt and the basic plumage of male Harlequin Ducks. Condor 99:83-90.
- Cottam, Clarence. 1939. Food habits of North American diving ducks. Pp. 80-86 in: Tech. Bull. No. 643. USDA, Washington, D.C.
- Coues, E. 1974. Birds of the Northwest. A hand-book of the ornithology of the region drained by the Missouri River and its tributaries. U.S. Geological Survey of the Territories Misc. Publ. 3. 791 pp.
- Crowley, D. W. 1994. Breeding habitat of Harlequin ducks in Prince William Sound, Alaska. P. 4 in: Proc. 2nd Ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Davis, C. V. 1961. A distributional study of the birds of Montana. Ph.D. dissertation. Oregon State University, Corvallis. 462 pp.
- Delacour, J. 1959. Waterfowl of the world. Vol. 3. Country Life Ltd., London. 270 pp.
- Delacour, J. and E. Mayr. 1945. The family Anatidae. Wilson Bull. 57(1):3-55.
- Dement'ev, G. P. and N. A. Gladkov, (eds.) 1967. Birds of the Soviet Union. Vol. 4. Translated from 1952 Russian edition, Israel Program for Scientific Translations. U.S. Department of the Interior and National Science Foundation, Washington, D.C.
- Derscheid, J. M. 1941. Notes sur certains Canards plongeurs. Gerfaut 31(3):87-99. (Abstr.)
- Diamond, S. and P. Finnegan. 1992. Harlequin duck ecology on Montana's Rocky Mountain Front. Unpublished report. Rocky Mountain District, Lewis and Clark National Forest, Choteau, MT. 45 pp.
- Diamond, S. and P. Finnegan. 1993. Harlequin duck ecology on Montana's Rocky Mountain Front. Unpublished report. Rocky Mountain District, Lewis and Clark National Forest, Choteau, MT. 45 pp.
- Dickinson, J. C., Jr. 1953. Report on the McCabe collection of British Columbian birds. Harvard University, Museum of

- Dow, D. D. 1964. Diving times of wintering water birds. *Auk* 81(4):556-558.
- Drolet, C. A. (ed). 1988. Report on a workshop on eider management held in Quebec city January 12-13, 1988. 53 pp.
- Dzinbal, K. A. 1982. Ecology of Harlequin ducks in Prince William Sound, Alaska, during summer. M.S. thesis. Ore. State Univ., Corvallis. 89 pp.
- Dzinbal, K. A. and R. L. Jarvis. 1982. Coastal feeding ecology of harlequin ducks in Prince William Sound, Alaska, during summer. Pp. 6-8 in: D. N. Nettleship et al. (eds.). *Marine birds: their feeding ecology and commercial fisheries relationships*. Canadian Wildlife Service Special Publication from Proc. Pacific Seabird Group Symposium, Seattle, Washington.
- Farman, L. M. and V. E. Miller. 1990. Results of 1990 surveys for harlequin ducks on the Kootenai and Lolo National Forests, Montana.
- Farman, L. M., D. L. Genter, and C. Jones. 1989. Results of the 1989 survey for harlequin ducks (*Histrionicus histrionicus*) on the Kootenai and Flathead National Forests, Montana. Unpublished report. 18 pp.
- Fay, F. H. 1960. The distribution of waterfowl to St. Lawrence Island, Alaska. *Wildfowl Trust Annual Report* 12:70-80.
- Fleischner, T. L. 1983. Natural history of Harlequin ducks wintering in northern Puget Sound. M.S. thesis. West. Washington Univ., Bellingham. 49 pp.
- Flint, V. E., R. L. Boehme, Y. V. Kostin, and A. A. Kuznetsov, (eds.). 1984. Harlequin duck. P. 50 in: *Birds of the USSR*. The Easton Press, Norwalk, Connecticut. 353 pp.
- Fournier, M. A., and R. G. Bromley. 1996. Status of the Harlequin Duck, *Histrionicus histrionicus*, in the western Northwest Territories. *Canadian Field-Nat.* 110:638-641.
- Gabrielson, I. N. and B. S. Wright. 1951. Notes on the birds of the Fort Chimo, Ungava District. *Canadian Field-Nat.* 65:127-140.
- Gabrielson, I. N. and F. C. Lincoln. 1959. *Birds of Alaska*. The Stackpole Co., Harrisburg, PA, and Wildl. Manage. Inst., Washington, D.C. 922 pp.
- Gabrielson, I. N. and S. G. Jewett. 1940. *Birds of Oregon*. Oregon State College, Corvallis, OR.
- Gaines, W. L. and R. E. Fitzner. 1987. Winter diet of Harlequin duck at Sequim Bay, Puget Sound, Washington. *Northwest Science* 61(4):213-215.
- Gangemi, J. T. 1991. Results of the 1991 survey for Harlequin Duck (*Histrionicus histrionicus*), distribution in the non-wilderness portion of the Flathead National Forest, Montana. Unpublished report for the MTNHP. 26 pp.
- Gardarsson, A. 1979. Waterfowl populations of Lake Myvatn and recent changes in numbers and food habits. *Oikos* 32:250-270.
- Gardarsson, A., G. M. Gislason, and A. Einarsson. 1988. Long term changes in the Lake Myvatn ecosystem. *Aqua Fennica* 18:2:125-135.
- Genter, D. L. 1992. Status of the Harlequin duck in Montana. P. 5 in: *Proc. Harlequin duck Symposium*, April 23-24, 1992, Moscow, Idaho. ID Dept. of Fish & Game, U.S. For. Serv. Intermtn. Res. Stat., ID Panhandle Nat. Forests, and NW Sect. of Wildl. Soc. 46 pp.

- Genter, D. L. 1993. Harlequin duck status report 1992. Montana. Pp. 31-34 in: Cassirer, E. F., et al., (eds.), Status of harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Genter, D. L. 1993. Whitewater wonder. Montana Outdoors 24(4): 2-7 July/August.
- Genter, D. L. and J. D. Reichel. 1994. Harlequin duck surveys in western Montana. 1994. P. 19 in: Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Gislason, G. M. 1985. The life cycle and production of *Simulium vittatum* Zett. in the River Laxa, North-East Iceland. International Association of Theoretical and Applied Limnology 22: 3281-3287.
- Gislason, G. M. 1994. River management in cold regions: a case study of the River Laxa, North Iceland. Pp. 464-483 in: P. Calow and G. E. Petts, (eds.), The Rivers Handbook--Hydrological and ecological principles. Blackwell Scientific Publications, Oxford.
- Gislason, G. M. and A. Gardarsson. 1988. Long term studies on *Simulium vittatum* Zett. (Diptera: Simuliidae) in the River Laxa, North Iceland, with particular reference to different methods used in assessing population changes. International Association of Theoretical and Applied Limnology 23: 2179-2188.
- Godfrey, W. E. 1986. The birds of Canada, revised edition. Distributed for National Museum of Natural Sciences, Ottawa Univ. Chicago Press. 596 pp.
- Goudie, R. I. 1984. Comparative ecology of Common eiders, black scoters, oldsquaws and harlequin ducks wintering in southeast Newfoundland. Thesis. Univ. of W. Ontario, London, Ontario, Canada.
- Goudie, R. I. 1988. Breeding distribution of harlequin ducks in northern Labrador. Atlantic Soc. of Fish and Wildl. Biologists. 4(2): 17-21.
- Goudie, R. I. 1989. Historical status of harlequin ducks wintering in eastern North America: a reappraisal. Wilson Bull. 101: 112-114.
- Goudie, R. I. 1991. The status of the Harlequin Duck (*Histrionicus histrionicus*) in eastern North America. Revised 15 Feb 1991. Committee on the status of endangered wildlife in Canada (COSEWIC), Ottawa, Ontario. 59 pp. + 4 appendices.
- Goudie, R. I. 1993. Harlequin duck status report. eastern Canada. Pp. 65-74 in: Cassirer, E. F., et al., (eds.), Status of Harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Goudie, R. I. and C. D. Ankney. 1988. Patterns of habitat use by sea ducks wintering in southeastern Newfoundland. Ornis Scandinavica 19: 249-256.
- Goudie, R. I. and C. D. Ankney. 1986. Body size, activity budgets, and diets of sea ducks wintering in Newfoundland. Ecology 67(6): 1475-1482.
- Goudie, R. I. and S. Brault. 1994. Lords and ladies: an exercise in sea duck modeling. Pp. 15-16 in: Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Goudie, R. I., S. Brault, B. Conant, A. V. Kondratyev, M. R. Petersen, K. Vermeer. 1994. The status of sea ducks in the North Pacific rim: toward their conservation and Management. Pg. 27-49 in: Transactions of the 59th North American Wildlife and Natural Resources Conference.
- Grinnell, J. and A. H. Miller. 1944. The distribution of the birds of California. Cooper Ornithological Club, Pacific Coast Avifauna, No. 27.

- Groves, C., Wallen, W. and F. Cassirer. 1990. Clown on the water. Idaho Wildlife 10(3):24-25.
- Gudmundsson, F. 1971. Straumendur (*Histrionicus histrionicus*) a Islande. ("The harlequin duck in Iceland") Naturufroedingurinn 41(1):1-28, (2):64-98. (English summary pp. 84-98).
- Hand, R. L. 1932. Notes on the occurrence of water and shore birds in the Lochsa region of Idaho. Condor 17:118-129.
- Hand, R. L. 1941. Birds of the St. Joe National Forest, Idaho. Condor 43:220-232.
- Hand, R. L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.
- Harju, H. 1980. Harlequin ducks in Wyoming. Wyoming Wildlife, Jan., pp. 16-17.
- Harlequin Duck Working Group. 1992. Proceedings of Harlequin Duck Symposium. held at the Northwest Section of the Wildlife Society Meeting, April 23-24, 1992, Moscow, Idaho. Idaho Department of Fish and Game. 45 pp.
- Harlequin Duck Working Group. 1993. Status of harlequin ducks (*Histrionicus histrionicus*) in North America. Report of the Harlequin Duck Working Group. 83 pp.
- Harlequin Duck Working Group. 1994. Proceedings of the second Harlequin Duck symposium. Harlequin Duck Working Group, Site 12, Box 15, RR 31, Galiano, B.C. V0N 1P0. 22 pp.
- Harrison, J. 1967. Drake harlequin escorting its family. Wildfowl Trust Ann. Report 18:155-156.
- Hendricks, P., and J. D. Reichel. 1998. Harlequin Duck research and monitoring in Montana: 1997. Montana Natural Heritage Program. Helena, MT. 28 pp.
- Hirsch, K. V. 1980. Winter ecology of sea ducks in the inland marine waters of Washington. M.S. thesis. University of Washington. 92 pp.
- Hoffmann, R. 1927. Birds of the Pacific states. Houghton Mifflin, Boston.
- Hunt, B. 1994. Behavioral ecology of breeding Harlequin ducks in Jasper National Park: time-activity budgets and the effects of human disturbance. P. 17 in: Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Hunt, B. and P. Clarkson. 1993. Preliminary status and distribution of Harlequin ducks on selected breeding ranges in the Canadian Rockies. Pp. 75-83 in: Cassirer, E. F., et al., (eds.), Status of Harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Hunt, W. A. 1993. Jasper National Park harlequin duck research project, 1992 pilot projects--interim results. Jasper Warden Service Biological Report Series, No. 1. Heritage Resource Conservation, Parks Canada, Box 10, Jasper, Alberta. 67 pp.
- Inglis, I. R., J. Lazarus, and R. Torrance. 1989. The pre-nesting behavior and time budget of the Harlequin duck *Histrionicus histrionicus*. Wildfowl 40:55-73.
- Irving, L. 1960. Birds of the Anaktuvuk Pass, Kobuk, and Old Crow. U.S. Natl. Mus. Bull. 217. 409 pp.
- Jewett, S. G. 1931. Nesting of the Pacific harlequin duck in Oregon. Condor 33:255.
- Jewett, S. G., W. P. Taylor, W. T. Shaw, and J. W. Aldrich. 1953. Birds of Washington State. University of Washington Press, Seattle. 767 pp.

- Johnsgard, P. A. 1962. Evolutionary trends in the behaviour and morphology of the Anatidae. Wildfowl Trust Ann. Rep. 13: 130-148.
- Johnsgard, P. A. 1965. Handbook of waterfowl behavior. Comstock Press, Ithaca, NY. 378 pp.
- Johnsgard, P. A. 1975. Waterfowl of North America. Indiana University Press, Bloomington. 304 pp.
- Johnson, D. D. 1991. Results of stream surveys for Harlequin ducks in the Gallatin and a section of the Custer National Forests, Montana. Unpublished report to the Montana Natural Heritage Program. 18 pp.
- Johnstone, S. T. 1970. Waterfowl eggs. Aviculture Mag. 76:52-55.
- Koskimies, J. and L. Lahti. 1964. Cold-hardiness of the newly hatched young in relation to ecology and distribution in ten species of European ducks. Auk 81(3): 281-307.
- Kuchel, C. R. 1977. Some aspects of the behavior and ecology of harlequin ducks breeding in Glacier National Park, Montana. M.S. thesis. Univ. of Montana, Missoula. 160 pp.
- Kurechi, M. and K. Yamada. 1984. Foods in the gizzard of a male harlequin duck. Tori 33: 78-80.
- Larrison, E. J., J. L. Tucker, and M. T. Jollie. 1967. Guide to Idaho birds. J. Idaho Acad. of Sci. 5: 1-220.
- Latta, S. C. 1993. Distribution and status of the Harlequin duck (*Histrionicus histrionicus*) in Oregon. Pp. 35-44 in: E. F. Cassirer et al., (eds.), Status of Harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Lee, D. N. B., and D. L. Genter. 1991. Results of harlequin duck (*Histrionicus histrionicus*) surveys in wilderness areas of the Flathead National Forest, Montana. Montana Natural Heritage Program. Helena, MT. 31 pp.
- Lokemoen, J. T. and D. E. Sharp. 1985. Assessment of nasal marker materials and designs used on dabbling ducks. Wildlife Society Bulletin 13: 53-56.
- Markum, D. and D. L. Genter. 1990. Preliminary report on the distribution and status of the harlequin duck, *Histrionicus histrionicus* on the Gallatin National Forest, Montana. Unpublished report for the Gallatin National Forest. Montana Natural Heritage Program. 21 pp.
- McEneaney, T. 1994. Status of the Harlequin duck in Yellowstone National Park, Wyoming. P. 5 in: Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- McEneaney, T. 1997. Harlequins: noble ducks of turbulent waters. Yellowstone Science 5(Spring): 2-7.
- Merriam, C. H. 1883. Breeding of the harlequin duck (*Histrionicus histrionicus*) in Newfoundland. Bull. Nuttall Ornithology Club 8: 200.
- Merrill, J. C. 1897. Notes on the birds of Fort Sherman, Idaho. Auk 14: 347-357.
- Meyer de Schauensee, R. 1984. The birds of China. Smithsonian Institution Press, Washington, D.C. 602 pp.
- Michael, C. W. and E. Michael. 1922. An adventure with a pair of harlequin ducks in the Yosemite Valley. Auk 39: 14-23.
- Miller, V. E. 1988. Harlequin ducks (*Histrionicus histrionicus*) 1988 results of field survey in west-central Montana. 18 pp.
- Miller, V. E. 1989. Field survey report, harlequin duck (*Histrionicus histrionicus*) lower Clark Fork River drainage, west-central Montana. Unpublished report. 47 pp.

- Mittelhauser, G. H. 1989. The ecology and distribution of the harlequin duck (*Histrionicus histrionicus*) wintering off Isle au Haut, Maine. B.A. thesis. College of the Atlantic. 69 pp.
- Mittelhauser, G. H. 1990. Survey of harlequin ducks along the Isle au Haut shoreline and adjacent offshore islands, 1989-90. Unpublished report, College of the Atlantic, Bar Harbor, Maine. 25 pp.
- Mittelhauser, G. H. and M. McCollough. 1993. Harlequin duck status report 1992. Maine. Pp. 55-59 in: Cassirer, E. F., et al., (eds.), Status of Harlequins ducks in North America. Harlequin Duck Working Group. 83 pp.
- Mittelhauser, G. and J. Hazen. 1990. Monitoring harlequins at Acadia. Park Sci. 10(1):18.
- Montana Department of Natural Resources and Conservation. 1980. Kootenai Falls wildlife monitoring study, first annual report for the period September 2, 1979 - September 1, 1980. Unpublished report. 40 pp.
- Montana Department of Natural Resources and Conservation. 1981. Kootenai Falls wildlife monitoring study, second annual report for the period September 2, 1980 - September 1, 1981. 33 pp.
- Montana Department of Natural Resources and Conservation. 1982. Kootenai Falls wildlife monitoring study, third annual report for the period September 2, 1981 - September 1, 1982. 36 pp.
- Montana Department of Natural Resources and Conservation. 1983. Kootenai Falls wildlife monitoring study, fourth annual report for the period September 1, 1982 - September 1, 1983. Unpublished report. 39 pp.
- Montevvecchi, W. A., et al. 1995. National recovery plan for the Harlequin Duck in eastern North America. Prepared by the Harlequin Duck (eastern North Am. pop.) Recovery Team for the Recovery of Nationally Endangered Wildlife Committee. 31 pp.
- Morneau, F. and R. Decarie. 1994. Status and distribution of Harlequin ducks in the Great Whale Watershed, Quebec. Pp. 6-7 in. Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994 Harlequin Duck Working Group. 22 pp. plus appendices.
- Murie, O. J. 1959. Fauna of the Aleutian Islands and Alaska Peninsula. U.S. Department of the Interior, Fish and Wildlife Service, North American Fauna, No. 61. 406 pp.
- Myres, M. T. 1959. The behavior of sea ducks and its value in the systematics of the tribes Mergini and Somaterini of the family Anatidae. Ph.D. dissertation. Univ. of British Columbia, Vancouver.
- Nakashima, D. 1987. Common eider banding project in east Ungava Bay: experimental capture by net. Rep. prepared for Serv. de la Faune, Quebec Reg., Ste-Foy, Quebec. 38 pp.
- Nelson, A. D. and A. C. Martin. 1953. Gamebird weights. Journal of Wildlife Management 17:36-42.
- Palmer, R. S. 1949. Maine birds. Bull. Museum Comparative Zoology (Harvard) 102:110-117.
- Palmer, R. S. 1976. Handbook of North American birds. Volume 3. Yale University Press, New Haven, Connecticut.
- Palmer, T. S. 1921. The harlequin duck in Montana. Condor 23:133.
- Parkes, K. C. and C. H. Nelson. 1976. A definite Colorado breeding record for the Harlequin Duck. Auk 93:846-847.
- Patten, S. 1993. Acute and sublethal effects of the Exxon Valdez oil spill on harlequins and other seaducks. Alaska Dept. Fish and Game, Anchorage, AK. 7 pp.

- Patten, S. 1993. Reproductive failure of harlequin ducks. *Alaska's Wildlife*. January/February. 14-15.
- Patten, S. M., Jr. 1994. Assessment of injury to Harlequin ducks from the Exxon Valdez oil spill. P. 8 in Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Pearse, T. 1942. Notes on the migration of ducks on the east coast of Vancouver Island, B.C. *Murrelet* 23(1): 14-16.
- Pearse, T. 1945. Mating of the Pacific Harlequin Duck. *Canadian Field-Naturalist* 59: 66-67.
- Pearse, T. 1946. Notes on changes in bird populations in the vicinity of Comox, Vancouver Island--1917 to 1944. *Murrelet* 27(1): 4-9.
- Peters, H. S. and T. D. Burleigh. 1951. The birds of Newfoundland. U.S. Fish and Wildlife Service. 431 pp.
- Phillips, J. C. 1986. A natural history of the ducks, vol. III. Dover Publications, Inc., NY. 383 pp.
- Pinel, H. W., W. W. Smith, and C. R. Wershler. 1991. Harlequin duck (*Histrionicus histrionicus*). Pp. 62-62 in Alberta Birds, 1971-1980, Vol. 1. Non-Passerines. Natural History Section, Provincial Museum of Alberta, Edmonton, Alberta.
- Pool, W. 1962. Feeding habits of the harlequin. *Wildfowl Trust Ann. Rep.* 13: 126-129.
- Portenko, L. A. 1981. *Histrionicus histrionicus pacificus* Brooks - Harlequin duck. Pp. 179-182 in Birds of the Chukchi Peninsula and Wrangel Island. Nauka Publishers, Leningrad. 446 pp.
- Reel, S. L., Schassberger, W. Ruediger. 1989. Caring for our natural community. Region 1 - Threatened, Endangered & Sensitive Species Program. U.S.D.A., Forest Serv., Missoula, MT. 334 pp.
- Reichel, J. D. 1996. Literature review and summary of research priorities for the Harlequin Duck. Montana Natural Heritage Program, Helena, MT. 37 pp.
- Reichel, J. D. and D. L. Genter. 1993. Harlequin duck surveys in western Montana for 1992. Montana Natural Heritage Program, Helena, MT. 67 pp., including appendices and maps.
- Reichel, J. D. and D. L. Genter. 1994. Harlequin duck surveys in western Montana. 1993. Montana Natural Heritage Program, Helena, MT. 87 pp.
- Reichel, J. D. and D. L. Genter. 1995. Harlequin duck surveys in western Montana. 1994. Montana Natural Heritage Program, Helena, MT. 58 pp.
- Reichel, J. D. and D. L. Genter. 1996. Harlequin duck surveys in western Montana. 1995. Montana Natural Heritage Program, Helena, MT. 107 pp.
- Reichel, J. D., D. L. Genter and D. P. Hendricks. 1997. Harlequin duck research and monitoring in Montana. 1996. Montana Natural Heritage Program. Helena, MT. 77 pp.
- Robertson, G. J., F. Cooke, R. I. Goudie, and W. S. Boyd. 1998a. The timing of pair formation in Harlequin Ducks. *Condor* 100: 551-555.
- Robertson, G. J., F. Cooke, R. I. Goudie, and W. S. Boyd. 1998b. Moults speed predicts pairing success in male Harlequin Ducks. *Animal Behavior* 55: 1677-1684.
- Rothe, J. C. 1994. Progress toward a statewide status report for Harlequin ducks in Alaska. P. 9 in Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.

- Salomonsen, F. 1950. Birds of Greenland Part I. Ejnar Munksgaard, Copenhagen. 158 pp.
- Sato, H. and I. Kominato. 1988. The breeding records and breeding ecology of the harlequin duck *Histrionicus histrionicus* along the Ichihazama River of Mt. Kurikoma, northern Honshu. *Strix* 7:175. [In Japanese with English abstract]
- Saunders, A. A. 1914. The birds of Teton and northern Lewis and Clark counties, Montana. *Condor* 16:124-144.
- Saunders, A. A. 1915. Harlequin duck in Glacier National Park, Montana. *Auk*. 32:225.
- Saunders, A. A. 1921. A distributional list of the birds of Montana. *Pac. Coast Avifauna* 14. 194 pp.
- Savard, J. P. L. and A. M. Breault. No date. Molting Harlequin ducks: capture technique, weights and movements. Unpublished draft. 16 pp.
- Schirato, G. 1993. A preliminary status report of Harlequin ducks in Washington: 1993. Pp. 45-48 in: Cassirer, E. F., et al., (eds.), Status of Harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Schirato, G. 1994. Population and distribution of Harlequin ducks in Washington State. P. 18 in: Proc. 2nd ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.
- Schirato, G. and F. Sharpe. 1992. Distribution and habitat use of Harlequin ducks in northwestern Washington. P. 4 in: Proc. Harlequin duck Symposium, April 23-24, 1992, Moscow, Idaho. ID Dept. of Fish & Game, U.S. For. Serv. Intermtn. Res. Stat., ID Panhandle Nat. Forests, NW sect. of Wildl. Soc. 46 pp.
- Smith, C. 1996. Banff National Park Harlequin duck research project: progress report - 1995 field season. Heritage Resource Conservation, Parks Canada, Banff, Alberta, Canada. 41 pp. plus appendices.
- Smith, C. 1999. Banff National Park Harlequin Duck research project – 1998 progress report. Unpublished technical report. Heritage Resource Conservation, Banff National Park, Alberta, Canada. 40 pp.
- Snyder, L. L. 1957. Arctic birds of Canada. Univ. of Toronto Press, Toronto.
- Soper, J. D. 1946. Ornithological results of the Baffin Island expeditions of 1928-1929 and 1930-1931, together with more recent records. *Auk* 63:1-24.
- Soper, J. D. 1954. Waterfowl and other ornithological investigations in Yukon Territory, Canada, 1950. *Can. Wildl. Serv. Wildl. Manage. Bull. (Ser. 2)* 7. 55 pp.
- Taylor, P. and J. E. Thompson 1990. Harlequin duck in Manitoba: an update. *Blue Jay* 48(2):98-103.
- Thompson, J., R. Goggans, P. Greenlee, and S. Dowlan. 1993. Abundance, distribution and habitat associations of the harlequin duck (*Histrionicus histrionicus*) in the Cascade Mountains, Oregon. Unpublished report prepared for cooperative agreement between the Oregon Department of Fish and Wildlife, Willamette National Forest, Mt. Hood National Forest, and the Bureau of Land Management, Salem District. 37 pp.
- Thompson, J., R. Goggans, P. Greenlee, and S. Dowlan. 1994. Abundance, distribution and habitat associations of the harlequin duck (*Histrionicus histrionicus*) in the Cascade Mountains, Oregon. [Abstract]. Proceedings Second Harlequin Duck Symposium, Hornby Island, B.C., 13-15 March 1994 2:1.
- Thompson, L. 1985. A harlequin romance. *Montana Outdoors* 16:21-25
- Titone, J. 1989. Biologists track reclusive harlequin duck. *The Spokane Chronicle*, 9 July 1989:1. Spokane, WA.



- Todd, W. E. C. 1963. Birds of the Labrador Peninsula and adjacent areas. Carnegie Mus. and Univ. of Toronto Press, Toronto. 819 pp.
- Tufts, R. W. 1961. The birds of Nova Scotia. Nova Scotia Mus., Halifax, N.S. 481 pp.
- Vermeer, K. 1983. Diet of the harlequin duck in the Strait of Georgia, British Columbia. Murrelet 64: 54-57.
- Vickery, P. D. 1988. Distribution and population status of harlequin ducks (*Histrionicus histrionicus*) wintering in eastern North America. Wilson Bull. 100: 119-126.
- Wade, J. M. 1881. Rare finds. Ornithologist 6(6): 44.
- Wallen, R. 1992. Annual variation in Harlequin duck population size, productivity and fidelity in Grand Teton National Park, Wyoming. P. 7 in Proc. Harlequin duck Symposium, April 23-24, 1992, Moscow, Idaho. ID Dept. of Fish & Game, U.S. For. Serv. Intermountain Res. Stat., ID Panhandle Nat. Forests, and NW Sect. of the Wildl. Soc. 46 pp.
- Wallen, R. 1992. Harlequin duck status report 1992. Wyoming. Pp. 49-54 in Cassirer, E. F., et al., (eds.), Status of Harlequin ducks in North America. Harlequin Duck Working Group. 83 pp.
- Wallen, R. L. 1987. Habitat utilization by Harlequin ducks in Grand Teton National Park. M.S. thesis. Montana State Univ., Bozeman. 67 pp.
- Wallen, R. L. 1987b. Annual brood survey for harlequin ducks in Grand Teton National Park. Grand Teton Nat. Pk., Resource Manage. 15 pp.
- Wallen, R. L. 1991. Annual variation in harlequin duck population size, productivity and fidelity to Grand Teton National Park. Off. of Science and Res. Mgt. Grand Teton National Park, WY. 7 pp.
- Wallen, R. L. and C. R. Groves. 1988. Status and distribution of harlequin ducks (*Histrionicus histrionicus*) in northern Idaho. Idaho Department of Fish and Game, Boise, ID. 34 pp.
- Wallen, R. L. and C. R. Groves. 1989. Distribution, breeding biology and nesting habitat of harlequin ducks (*Histrionicus histrionicus*) in northern Idaho. Report on Challenge Cost Share Project. 40 pp.
- Warren, E. R. 1914. Harlequin duck in Glacier National Park, Montana. Auk. 31: 535.
- Wild Bird Society of Japan. 1989. Sea ducks: harlequin duck. P. 56 in Sonobe, K. and J. W. Robinson, (eds.), A field guide to the birds of Japan. Kodansha International Ltd., Tokyo, New York, San Francisco.
- Yocum, C. F. 1951. Waterfowl and their food plants in Washington. Univ. of Washington Press, Seattle. 272 pp.
- Zwiefelhofer, D. C. 1994. Status of Harlequin ducks (*Histrionicus histrionicus*) on Kodiak Island, Alaska. P. 10 in Proc. 2nd Ann. Harlequin duck symposium, March 13-15, 1994. Harlequin Duck Working Group. 22 pp. plus appendices.

Appendix 1. Results of Harlequin Duck surveys in 1998 on the Lower Clark Fork tributaries. M and F in comments refer to resighted previously-banded males and females (usually blue color bands), juvenile age classes are as described in the methods.

STREAM	DATE	PAIRS	MALES	FEMALES	BROODS	YOUNG	COMMENTS
Rock Creek	9 May	1	1	--	--	--	F:VA
	19 May	1	1	1	--	--	F:VA
	5 Jul	--	--	--	--	--	--
	14 Jul	--	--	2	2	11	7 IIA, 4 IIB juvs.
	31 Jul	--	--	1	1	2	A third brood: 2 IIA juvs.
Marten Creek	3/4 Aug	--	--	--	--	--	older broods fledged
	5 Aug	--	--	1	1	2	F:TC, early IIB juvs.
	10 May	7	2	--	--	--	M:S1, S8, S9, F:SE
Swamp Creek	20 May	2	3	--	--	--	F:SE
	4/5 Jul	--	--	2	1	4	4 IIA juvs.
	17 Jul	--	--	2	2	11	F:SA; 6 IIC, 5 III juvs.
	1 Aug	--	--	--	--	--	Older brood fledged
	4 Aug	--	--	1	1	5	F:SC; 5 flying juvs.
Vermilion River	13 May	--	--	--	--	--	No ducks
	22 May	--	--	--	--	--	No ducks
	6 Jul	--	--	--	--	--	No ducks
	15 Jul	--	--	--	--	--	No ducks
	31 Jul	--	--	--	--	--	No ducks
Vermilion River	12 May	2	--	--	--	--	F:X1
	21 May	--	1	--	--	--	--
	7 Jul	--	--	1	1	3	3 IIA juvs.
	16/17 Jul	--	--	2	2	7	4 IB, 3 IIB juvs.
	2 Aug	--	--	--	--	--	Mouth to Grouse Creek
	3 Aug	--	--	2	2	6	F:925-09393; 4IIA, 2IIC juvs.

Appendix 2. Previously banded Harlequin Ducks observed in northwestern Montana during 1998 MTNHP surveys. See Ashley (in prep.) for all 1998 Glacier National Park observations.

Stream where Resighted	Band No. *	Sex	Date first Resighted	Stream where Banded	Date Banded	Age when Banded
Rock Cr	VA	F	9 May	Swamp	25 Jul 95	Juv.
Rock Cr	TC	F	5 Aug	Rock	26 Jul 95	Juv.
Marten Cr.	S1	M	10 May	Marten	26 Apr 95	Adult
Marten Cr	S8	M	10 May	Marten	10 May 96	Adult
Marten Cr	S9	M	10 May	Marten	10 May 96	Adult
Marten Cr	SE	F	10 May	Marten	28 Jul 95	Juv.
Marten Cr	SA	F	17 Jul	Marten	28 Jul 95	Juv.
Marten Cr	SC	F	4 Aug	Marten	28 Jul 95	Juv.
Vermilion R	X1	F	12 May	Vermilion	29 Jul 95	Adult
Vermilion R	X2 <sup>1</sup>	F	3 Aug	Vermilion	31 Jul 96	Adult
Trail Cr	F7	F	15 May	Trail	4 May 96	Adult
Trail Cr	J8	F	7 Aug	Trail	1 Aug 96	Adult
McDonald Cr	G3	F	6 Aug	McDonald	12 Aug 97	Adult
McDonald Cr	B5	F	19 Aug	McDonald	4 May 95	Adult
McDonald Cr	B7 <sup>2</sup>	F	19 Aug	McDonald	2 Sep 92	Adult

\* Blue color band, unless indicated otherwise.

<sup>1</sup> Aluminum band 925-09393. Color band added in 1998.

<sup>2</sup> Original aluminum band 755-76054, new band 925-72518. Color band added in 1998.

Appendix 3. Annual productivity of Harlequin Ducks during 1992-1998 on the four Lower Clark Fork streams, Sanders County, Montana.

STREAM	1992	1993	1994	1995	1996	1997	1998
<b>Marten Creek</b>							
No. adult females	4	5	5	5	5	3	7
No. broods	4	2	0	2	2	1	2
Broods/female	1.00	0.40	0	0.40	0.40	0.33	0.29
Young/brood	3.25	4.00	0.00	5.5	2.5	6.00	5.5
<b>Rock Creek</b>							
No. adult females	0	2 (?)	4	1(?)	3	1	3
No. broods	0	2	1	1	0	1	3
Broods/female	0	1.00	0.25	1.00	0.00	1.00	1.00
Young/brood	0.00	3.00	3.00	5.00	0.00	6.00	4.33
<b>Swamp Creek</b>							
No. adult females	0	1	0	3	2	0	0
No. broods	0	1	0	2	2	0	0
Broods/female	0.00	1.00	0.00	0.67	1.00	0.00	0.00
Young/brood	0.00	1.00	0.00	5.00	2.50	0.00	0.00
<b>Vermilion River</b>							
No. adult females	1	2	2	2	2	1	3
No. broods	0	2	1	2	1	0	2
Broods/female	0.00	1.00	0.50	1.00	0.50	0.00	0.67
Young/brood	0.00	3.00	6.00	2.50	7.00	0.00	3.00
<b>Combined Streams</b>							
No. females	5	10	11	11	12	5	12
No. broods	4	7	2	7	5	2	7
Broods/female	0.80	0.70	0.18	0.73	0.42	0.40	0.58
Young/brood	3.25	3.00	4.50	4.43	3.40	6.00	4.29



